TEXT SEARCHABLE DOCUMENT - 2010

Data Evaluation Report on the Acute Toxicity of DPX-MAT28 Technical (Aminocyclopyrachlor Acid) to Fish (Lepomis macrochirus)

PIVIKA Subillissi	on Number {	}		EPA MRID	Number 4/560124	
Data Requireme	E C E	PMRA Data Code EPA DP Barcode DECD Data Point EPA MRID EPA Guideline	{} D358148 {} 47560124 850.1075			-
Test material: Common name: Chemical name:	6-Amin CAS No. 858956-	lor acid Technical 5-chloro-2-cyclopro o-5-chloro-2-cyclop o-5-chloro-2-cyclop	opylpyrimidine-4 oropyl-4-pyrimid oropyl-pyrimidin			
•	er: John Marton Cambridge Environ	imental, Inc.	Signa Date	ature: : 07/01/09	Martin	
•	ver: Teri S. Myers Cambridge Enviro	onmental, Inc.	Signa Date	ature: : 07/01/09 ature: : 07/20/09 ature: : 10/06/09	i'S Mym	
Secondary Revi EPA/OPP/EFEI	ewer: Anita Ullagad D/ERB1	ldi	Signa Date	ature: 5777 : 10/06/09	~	
Reference/Subm	nission No.: {	}				
Company Code Active Code Use Site Catego EPA PC Code	{} [For PMRA]				
Date Evaluation	Completed: 10/06/	/09	*			

<u>CITATION</u>: Gallagher, S.P., T.Z. Kendall and H.O. Krueger. 2007. DPX-MAT28 Technical: A 96-Hour Static Acute Toxicity Test with the Bluegill (*Lepomis macrochirus*). Unpublished study performed by Wildlife International, Ltd., Easton, MD. Laboratory report number 112A-214. Study sponsored by E.I. du Pont de Nemours and Company, Wilmington, DE. Study completed November 29, 2007.

<u>DISCLAIMER</u>: This document provides guidance for EPA and PMRA reviewers on how to complete a data evaluation record after reviewing a scientific study concerning the acute toxicity of a pesticide to fish. It is not intended to prescribe conditions to any external party for conducting this study nor to establish absolute criteria regarding the assessment of whether the study is scientifically sound and whether the study satisfies any applicable data requirements. Reviewers are expected to review and to determine for each study, on a case-by-case basis, whether it is scientifically sound and provides sufficient information to satisfy applicable data requirements. Studies that fail to meet any of the conditions may be accepted, if appropriate; similarly, studies that meet all of the conditions may be rejected, if appropriate. In sum, the reviewer is to take into account the totality of factors related to the test methodology and results in determining the acceptability of the study.



PMRA Submission Number {......}

EPA MRID Number 47560124

EXECUTIVE SUMMARY:

In a 96-h acute toxicity study, bluegill sunfish (*Lepomis macrochirus*) were exposed to DPX-MAT28 (Aminocyclopyrachlor Acid) Technical at nominal concentrations of 0 (negative control), 7.5, 15, 30, 60 and 120 mg ai/L under static conditions; mean-measured concentrations were <0.0143 (<LOD; control), 7.5, 15, 29, 60 and 120 mg ai/L. The 96-h LC $_{50}$ was >120 mg ai/L. The EC $_{50}$ and NOAEC values were >120 and 120 mg ai/L, respectively, based on a complete lack of mortality and sub-lethal effects. Based on the results of this study, DPX-MAT28 Technical would be classified as practically non-toxic to *Lepomis macrochirus* in accordance with the classification system of the U.S. EPA.

This toxicity study is scientifically sound and classified as acceptable. It satisfies the guideline requirement for an acute freshwater fish toxicity study.

Results Synopsis

Test Organism Size/Age(mean weight or length): mean wet weight- 0.09 (0.05-0.13) g; mean total length- 2.1 (1.9-2.3) cm

Test Type (Flow-through, Static, Static Renewal): Static

LC₅₀: >120 mg ai/L

95% C.I.: N/A

NOAEC: 120 mg ai/L

Probit Slope: N/A

EC₅₀: >120 mg ai/L

95% C.I.: N/A

Endpoint(s) Affected: None

PMRA Submission Number {......}

EPA MRID Number 47560124

I. MATERIALS AND METHODS

GUIDELINE FOLLOWED:

This study was conducted following guidelines outlined in the OECD Guideline for Testing of Chemicals, 203, Fish Acute Toxicity Test; and U.S. Environmental Protection Agency Series 850- Ecological Effects Test Guidelines, OPPTS Number 850.1075, Fish Acute Toxicity Test, Freshwater and Marine (draft). The following deviations from OPPTS 850.1075 were noted:

1. The TOC and particulate matter of the dilution water were not reported.

2. While one replicate of ten fish per test level is acceptable, guidelines indicate a preference of two replicates per treatment with 10 fish each to improve statistical power by reducing variability unrelated to treatment.

These deviations does not affect the scientific soundness or acceptability of the study.

COMPLIANCE:

Signed and dated No Data Confidentiality, GLP and Quality Assurance statements were provided. This study was conducted in compliance with the U.S. EPA Good Laboratory Practice Standards (40 CFR Parts 160 and 792) that are consistent with: OECD Principles of Good Laboratory Practice (ENV/MC/CHEM (98) 17); and Japan MAFF (11 NohSan, Notification No. 6283, Agricultural Production Bureau, 1 October 1999), with the exception that the periodic analyses of well water for potential contaminants were performed using a certified laboratory and standard U.S. EPA analytical methods.

A. MATERIALS:

1. Test material

DPX-MAT28 Technical (Aminocyclopyrachlor)

Description:

Solid

Lot No./Batch No.:

Not Provided

Purity:

92.2%

Stability of compound

under test conditions:

Stable. Mean-measured concentrations ranged from 96.7 to 100% of

nominal.

(OECD recommends water solubility, stability in water and light, pKa, Pow, and vapor pressure of test compound)

Storage conditions of

test chemicals:

Stored at ambient temperature.

Physicochemical properties of DPX-MAT28 Technical.

Values	Comments
Not Reported	

PMRA Submission Number {......}

EPA MRID Number 47560124

Parameter	Values	Comments
Vapor pressure	Not Reported	
UV absorption	Not Reported	
pKa	Not Reported	
Kow	Not Reported	

2. Test organism:

Species:

Bluegill Sunfish (Lepomis macrochirus) EPA recommends a cold water species

(preferably rainbow trout Oncorhynchus mykiss) and a warm water species

(preferably bluegill sunfish Lepomis macrochirus). OECD recommends choice of

species at discretion of testing laboratory.

Age at test initiation:

Juvenile

Weight at study initiation: Length at study initiation: mean wet weight- 0.09 (0.05-0.13) g EPA recommends: mean 0.5 - 5 g. mean total length- 2.1 (1.9-2.3) cm EPA recommends: Longest not > 2x shortest; OECD recommends 2.0 \forall 1.0 cm for bluegill and 5.0 \forall 1.0 cm for

rainbow trout

Source:

Osage Catfisheries, Inc., Osage Beach, Missouri

EPA recommends that all organisms be from the same source

B. STUDY DESIGN:

1. Experimental Conditions

a. Range-finding study: The nominal concentrations were selected based upon exploratory range-finding results and in consultation with the study sponsor; no details pertaining to the range-finding trial was provided.

b. Definitive Study

Table 1: Experimental Parameters

Parameter	Details	Remarks Criteria		
I al amoto	Details			
Acclimation				
Period:	At least 14 days	The recommended acclimation period is a minimum of 14 days; OECD guideline		
Conditions: (same as test or not)	Same as test	recommends a minimum of 12 days. Pretest mortality should be < 3% 48 h.		
Feeding:	Fish were fed daily with a commercially-prepared diet (Ziegler Brothers, Inc., Gardners, Pennsylvania); fish were not fed for at least two days prior to testing.	prior to testing. OECD pretest mortality criteria: $>10\%$ = rejection of entire batch; ≥ 5 and $\leq 10\%$ = continued acclimation for 7 days; $<5\%$ = acceptable.		
Health: (any mortality observed)	No signs of stress were observed 14 days prior to testing and no mortality occurred during the 7 days prior to testing.			

PMRA Submission Number {.........}

EPA MRID Number 47560124

Parameter	Details	Remarks Criteria		
Duration of the test	96 hours			
		The recommended test duration is 96 hours.		
Test condition				
Static/flow-through	Static	A reproducible supply of toxicant is recommended. Consistent flow rate is		
Type of dilution system - for flow-through method.	N/A	usually 5-10 vol/24 hours, meter systems should be calibrated before and after study and checked twice daily during test period.		
Renewal rate for static renewal	N/A			
Aeration, if any	None reported			
		Aeration is not recommended; OECD guideline recommends aeration. If aeration is necessary, test solutions must be analyzed periodically to verify exposure.		
Test vessel		The state of the s		
Material: (glass/stainless steel)	Stainless steel	Test vessel size is usually 19 L (5 gal) or 30 x 60 x 30 cm.		
Size:	54 L	Fill volume is usually 15-30 L of solution.		
Fill volume:	40 L			
Source of dilution water Quality:	Water was obtained from an on-site well. Water was passed through a sand filter, pumped into a storage tank and aerated. Prior to use, water was filtered to 0.45 µm to remove fine particles.	Recommended source of dilution water is soft, reconstituted water or water from a natural source. EPA does not recommend the use of dechlorinated tap water; however, its use may be supportable if the biological responses for the organisms and chemical analyses of residual chlorine meet conditions in the Agency=s 850.1010 guidelines for dilution water (http://www.epa.gov/opptsfrs/OPPTS_H armonized/850_Ecological_Effects_Test_Guidelines/Draft/850.1010.pdf) Dilution water should be intensely aerated before the study. OECD permits dechlorinated tap water.		

PMRA Submission Number {......}

EPA MRID Number 47560124

Parameter	Details	Remarks		
1 ai aineel	Domin	Criteria		
Water parameters:		Specific Conductance= 320 μmhos/cm		
Hardness	138 mg/L as CaCO ₃ (0-hrs)	Alkalinity= 180 mg/L as CaCO ₃		
pH	7.8-8.5	<u>Hardness:</u> EPA recommends 40 - 48 mg/L as		
Dissolved oxygen	≥7.4 mg/L (≥85% saturation)	CaCO ₃ (OECD recommends 10 - 250 mg/L)		
Total Organic carbon	Not Reported	<u>pH</u> : EPA recommends 7.2 - 7.6; 8.0-8.3 for		
Particulate Matter	Not Reported	marine-stenohaline fishes, 7.7-8.0 for estuarine-euryhaline fishes, monthly		
Metals	See Reviewer's Comments	range < 0.8); (OECD recommends pH 6.0 - 8.5) Dissolved Oxygen:		
Pesticides	None Detected	EPA recommends: Static: 360% during first 48 hrs and 340% during second 48		
Chlorine	4.8 mg/L (Chloride)	hrs; flow-through: \$\frac{1}{2}60\%; (OECD) guideline recommends at least 80\%		
Temperature	21.5-22.8°C	saturation value). <u>Temperature</u> :		
{Salinity for marine or estuarine species}	N/A	EPA recommends 12 EC for coldwater species, 17 or 22 EC for warmwater species, and 22 ± 1 EC for		
Intervals of water quality measurement	Temperature, DO and pH were measured in every test vessel at test	estuarine/marine organisms. (OECD recommends 21 - 25°C for bluegill and 13 - 17°C for rainbow trout).		
	initiation and every 24 hours thereafter. Temperature was also measured continuously in the	Salinity: EPA recommends 30-34% (parts per thousand) for marine, 10-17% for		
	negative control. Hardness, alkalinity and conductance were	estuarine fish, weekly range < 6‰.		
	measured at test initiation.	Water quality should be measured at beginning of test and every 48 hours.		
Number of replicates/groups:				
control:		Recommended number of replicates		
solvent control:	N/A	include a control and five treatment		
treated ones:		levels with 2 replicates per level. Each concentration should be 60% of the next highest concentration, concentrations should be in a geometric series.		
Number of organisms per replicate				
/groups: control: solvent control:	10 N/A	Number of organisms per replicate should be 310/concentration; OECD		
treated ones:	10	guideline recommends at least 7 fish/concentration.		

Data Evaluation Report on the Acute Toxicity of DPX-MAT28 Technical (Aminocyclopyrachlor Acid) to Fish (Lepomis macrochirus) PMRA Submission Number {........}

EPA MRID Number 47560124

Parameter	Details	Remarks		
~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~		Criteria		
Biomass loading rate	0.02 g fish/L			
		Recommended static conditions are #0.8 g/L at #17EC and #0.5 g/L at > 17EC. Recommended flow-through conditions are #1 g/L/day. OECD recommends a maximum of 1 g fish/L for static and semi-static, while higher rates are recommended for flow-through.		
Test concentrations: nominal:	0 (negative control), 7.5, 15, 30, 60 and 120 mg ai/L			
measured:	<0.0143 (<lod; 120="" 15,="" 29,="" 60="" 7.5,="" ai="" and="" control),="" l<="" mg="" td=""><td></td></lod;>			
Solvent (type, percentage, if used)	N/A; a solvent was not used			
		The solvent should not exceed 0.5 ml/L for static tests or 0.1 ml/L for flow-through tests; OECD recommends that the solvent not exceed 100 mg/L.		
Lighting	16L:8D with a 30 minute transition period of low-light intensity	Light intensity at test initiation was 120 lux at the surface of the water of one representative test chamber.		
		The recommended photo period is 16 hours of light and 8 hours of dark with a 15-30 minute transition period. OECD recommends a photo period of 12-16 hours.		
Feeding	Fish were not fed during the study			
		Fish should not feed during the study.		
Recovery of chemical Frequency of determination Level of quantization Level of detection	0 and 96 hours 4.00 mg ai/L 0.0143 mg ai/L			
Positive control {if used, indicate the chemical and concentrations}	N/A; a positive control was not used			
Other parameters, if any	None reported			

PMRA Submission Number {......}

EPA MRID Number 47560124

2. Observations:

Table 2: Observations

Parameter	Details	Remarks Criteria		
Tarameter				
Parameters measured including the sublethal effects/toxicity symptoms	-mortality -sub-lethal effects			
Observation intervals	Observations were made after 4, 24,			
	48, 72 and 96 hours of exposure	Observation intervals should be a minimum of every 24 hours.		
Were raw data included?	Yes			
Other observations, if any	None reported			

II. RESULTS AND DISCUSSION:

A. MORTALITY:

No mortalities occurred in the control or treatment groups resulting in NOAEC and LC_{50} values of 120 and >120 mg ai/L, respectively.

PMRA Submission Number {......}

EPA MRID Number 47560124

Table 3: Effect of DPX-MAT28 Technical on Mortality of Lepomis macrochirus.

Mean-Measured and	No. of	Observation Period					
(Nominal) Concentrations	Fish at Start of Study	Day 1		Day 3		Day 4	
mg ai/L		No Dead	% Mortality	No Dead	% Mortality	No Dead	% Mortality
Negative Control	10	0	0	0	0	0	0
7.5 (7.5)	10	0	0	0	0	0	0
15 (15)	10	0	0	0	0	0	0
29 (30)	10	0	0	0	0	0	0
60 (60)	10	0	0	0	0	0	0
120 (120)	10	0	0	0	. 0.	. 0	0
NOAEC		-	* .	120 mg	ai/L		
LC ₅₀	>120 mg ai/L						
Positive control, if used mortality: LC ₅₀ :	N/A	N/A	N/A	N/A	N/A	N/A	N/A

N/A- Not Applicable

B. NON-LETHAL TOXICITY ENDPOINTS:

All control and treatment fish appeared normal and healthy throughout the definitive exposure period.

PMRA Submission Number {......}

EPA MRID Number 47560124

Table 4: Sub-lethal Effect of DPX-MAT28 Technical on Lepomis macrochirus.

Mean-Measured and	Observation Period					
(Nominal) Concentrations	Endpoints- Day 1	Endpoints- Day 3	Endpoints- Day 4 % Affected			
mg ai/L	% Affected	% Affected				
Negative Control	A.N.	A.N.	A.N.			
7.5 (7.5)	A.N.	A.N.	A.N.			
15 (15)	A.N.	A.N.	A.N.			
29 (30)	A.N.	A.N.	A.N.			
60 (60)	A.N.	A.N.	A.N.			
120 (120)	A.N.	A.N.	A.N.			
NOAEC		120 mg ai/L				
LOAEC		>120 mg ai/L				
EC ₅₀		>120 mg ai/L				
Positive control, if used % sublethal effect: EC ₅₀ :	Ŋ/A	N/A	N/A			

A.N.- All fish appeared normal and healthy

N/A- Not applicable

C. REPORTED STATISTICS:

All toxicity values were visually determined due to a lack of mortality.

D. VERIFICATION OF STATISTICAL RESULTS:

Statistical Method(s): The complete lack of sub-lethal effects and mortality precluded the statistical analysis of the data. Therefore, all toxicity values were visually determined based on the mean-measured concentrations.

LC₅₀: >120 mg ai/L

95% C.I.: N/A

NOAEC: 120 mg ai/L

Probit Slope: N/A

95% C.I.: N/A

E. STUDY DEFICIENCIES:

There were no study deficiencies.

F. REVIEWER=S COMMENTS:

The reviewer's results were identical to those of the study authors.

The results from the periodic screening analysis of the dilution water indicated the presence of the following elements: calcium (34.9 mg/L), chloride (4.8 mg/L), fluoride (1.2 mg/L), magnesium (12.8 mg/L), potassium (7.55 Page 10 of 11

PMRA Submission Number {......}

EPA MRID Number 47560124

mg/L) and sodium (19 mg/L).

The in-life portion of the definitive toxicity test was conducted from August 10 to August 14, 2007.

G. CONCLUSIONS:

This study is scientifically sound and classified as acceptable. The NOAEC and LC₅₀ values were 120 and >120 mg ai/L, respectively. Based on the results of this study, DPX-MAT28 Technical would be classified as practically nontoxic to *Lepomis macrochirus* in accordance with the classification system of the U.S. EPA.

 LC_{50} : >120 mg ai/L

95% C.I.: N/A

NOAEC: 120 mg ai/L

Probit Slope: N/A

95% C.I.: N/A

III. REFERENCES:

Organization for Economic Cooperation and Development. 1993. Guideline 203: Fish, Acute Toxicity Test. OECD Guideline for Testing of Chemicals. Updated Guideline adopted on 17 July 1992.

U.S. Environmental Protection Agency. 1996. OPPTS Number 850.1075: Fish Acute Toxicity Test, Freshwater and Marine. Series 850- Ecological Effects Test Guidelines (draft).

APHA, AWWA, WPCF. 1998. Standard Methods for the Examination of Water and Wastewater. 20th Edition. American Public Health Association. American Water Works Association. Water Pollution Control Federation, New York.

